

Treatment Protocols

Date: 07/01/2023

Abdominal Pain - Pediatric

Policy #9010P

Stable Blood pressure appropriate for age	Unstable Systolic blood pressure low for age, and/or signs of poor perfusion
Pediatric BLS Standing Orders	
<ul style="list-style-type: none"> • Universal Patient Protocol • In females, establish last menstrual period if patient has started her menses • Encourage immediate transport 	<ul style="list-style-type: none"> • Universal Patient Protocol • In females, establish last menstrual period if patient has started her menses • Encourage immediate transport
Pediatric LALS Standing Order Protocol	
<ul style="list-style-type: none"> • Establish IV PRN 	<ul style="list-style-type: none"> • Establish IV • 10-20 mL/kg NS IV bolus; titrated to age-appropriate systolic BP MR x1 • For persistent hypotension, see Shock Protocol
Pediatric ALS Standing Order Protocol	
<ul style="list-style-type: none"> • Establish IV/IO • Ondansetron 0.1 mg/kg, max 4 mg - IV/IO/IM/ODT PRN MR x1 • Pain Management Protocol PRN 	<ul style="list-style-type: none"> • Establish IV/IO • 10-20 mL/kg NS IV bolus; titrated to age-appropriate systolic BP MR x1 • Ondansetron 0.1 mg/kg, max 4 mg - IV/IO/IM/ODT PRN MR x1 • Pain Management Protocol PRN • For persistent hypotension, see Shock Protocol
Pediatric Base Hospital Orders	
<ul style="list-style-type: none"> • BH – repeat Ondansetron x 1 dose 	<ul style="list-style-type: none"> • BH - Repeat Ondansetron x 1 dose • BH – Repeat NS bolus x 1
Notes	
<ul style="list-style-type: none"> • In pediatrics (and less frequently adults), abdominal pain can be anaphylaxis. Consider epinephrine IM if clinical history and exam are consistent with anaphylaxis. 	

APPROVED:

Signature on File

Katherine Staats, M.D. FACEP

EMS Medical Director