

Treatment Protocols**Date: 07/01/2023****Hemodialysis Related Emergency/Suspected Hyperkalemia****Policy #9130A****Adult BLS Standing Orders**

- **Universal Patient Protocol**
- Give oxygen and/or ventilate PRN per **Airway Policy**
- Monitor O₂ saturation
- Glucose testing PRN
- Capnography
- For bleeding dialysis fistulas or shunts, refer to **Hemorrhage Control Policy**
- For bradycardia or tachycardia due to suspected electrolyte abnormalities, go to **Dysrhythmia Protocols**

Adult LALS Standing Orders

- Establish IV
- Capnography

Hypotension

- 250 mL NS IV MR to a max of 1,000 mL to maintain a SBP of ≥ 90 mmHg if patient is without rales and there is no evidence of heart failure

For Immediate Definitive Therapy Only:

- Establish IV in arm that does not have graft/AV fistula if possible, see **Policy on Pre-Existing Vascular Access Devices**

Adult ALS Standing Orders

- Monitor EKG
- Establish IV/IO
- Capnography

Hypotension

- 250 mL NS IV MR to a max of 1,000 mL to maintain a SBP of ≥ 90 mmHg if patient is without rales and there is no evidence of heart failure

For Immediate Definitive Therapy Only

- In life threatening conditions, and unable to obtain IO or vascular access, access graft/AV fistula, see **Policy on Pre-Existing Vascular Access Devices**

Fluid Overload with Rales

- Treat per **Respiratory Distress Policy**

Suspected Hyperkalemia (widened QRS complex or peaked T-waves)

- Obtain 12-Lead EKG

If abnormal - peaked t waves +/- widened QRS complex with symptoms of chest pain, shortness of breath, lightheadedness, weakness, or hypotension)

- Calcium Chloride 1,000 mg IV/IO
- Sodium Bicarbonate 50 mEq IV/IO
- Continuous Albuterol 5 mg via nebulizer

Peaked T-Waves are Defined as: > 5 mm in limb leads and/or > 10 mm in chest leads

APPROVED:

Treatment Protocols

Date: 07/01/2023

Hemodialysis Related Emergency/Suspected Hyperkalemia

Policy #9130A

Signature on File

Katherine Staats, M.D. FACEP

EMS Medical Director

Draft