

**Treatment Protocols**

**Date: 07/01/2023**

***Respiratory Distress or Failure - Pediatric***

**Policy #9170P**

<b>Stable</b> Systolic blood pressure appropriate for age	<b>Unstable</b> Systolic blood pressure low for age, and/or signs of poor perfusion
<b>Pediatric BLS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• <b>Universal Patient Protocol</b></li> <li>• Ensure patent airway, give oxygen and/or ventilate PRN per <b>Airway Policy</b></li> <li>• Maintain O2 saturation &gt; 95%</li> <li>• Capnography</li> <li>• Suction aggressively as needed</li> <li>• For adult-sized pediatric patients, can consider <b>NIPPV</b> – see <b>NIPPV procedure</b></li> <li>• Consider early <b>BHP contact</b></li> </ul> <p><b><u>RESPIRATORY DISTRESS WITH SUSPECTED BRONCHOSPASM</u></b></p> <ul style="list-style-type: none"> <li>• May assist patient with prescribed albuterol inhaler</li> </ul> <p><b><u>SUSPECTED ACUTE STRESSOR/HYPERVENTILATION SYNDROME</u></b></p> <ul style="list-style-type: none"> <li>• Remove from any causative environment</li> <li>• Coaching / reassurance</li> <li>• Do not utilize bag or mask rebreathing</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Universal Patient Protocol</b></li> <li>• Ensure patent airway, give oxygen and/or ventilate PRN per <b>Airway Policy</b></li> <li>• Maintain O2 saturation &gt; 95%</li> <li>• Capnography</li> <li>• Suction aggressively as needed</li> <li>• For adult-sized pediatric patients, can consider <b>NIPPV</b> – see <b>NIPPV procedure</b></li> <li>• Consider early <b>BHP contact</b></li> </ul> <p><b><u>RESPIRATORY DISTRESS WITH SUSPECTED BRONCHOSPASM</u></b></p> <ul style="list-style-type: none"> <li>• May assist patient with prescribed albuterol inhaler</li> </ul> <p><b><u>SUSPECTED ACUTE STRESSOR/HYPERVENTILATION SYNDROME</u></b></p> <ul style="list-style-type: none"> <li>• Remove from any causative environment</li> <li>• Coaching / reassurance</li> <li>• Do not utilize bag or mask rebreathing</li> </ul>
<b>Pediatric LALS Standing Order Protocol</b>	
<ul style="list-style-type: none"> <li>• Establish IV access PRN</li> <li>• Capnography</li> </ul> <p><b><u>SUSPECTED BRONCHOSPASM</u></b> (Suspected asthma)</p> <ul style="list-style-type: none"> <li>• Albuterol via nebulizer per ped dosing chart</li> </ul>	<ul style="list-style-type: none"> <li>• Establish IV</li> <li>• Capnography</li> </ul> <p><b><u>HYPOTENSION IF CARDIAC CAUSE NOT SUSPECTED</u></b></p> <ul style="list-style-type: none"> <li>• 10-20 mL/kg NS IV bolus; titrated to age-appropriate systolic BP MR x1, if patient is without rales and there is no evidence of heart failure</li> </ul> <p><b><u>SUSPECTED BRONCHOSPASM</u></b> (Suspected Asthma)</p> <ul style="list-style-type: none"> <li>• Albuterol via nebulizer per pediatric dosing chart</li> </ul> <p><b>If severe respiratory distress with bronchospasm or inadequate response to Albuterol/Atrovent, consider</b></p> <ul style="list-style-type: none"> <li>• Epinephrine 1:1,000 per drug chart IM SO. MR x 2 q5minutes</li> </ul> <p><b>Respiratory Distress with stridor at rest</b></p> <ul style="list-style-type: none"> <li>• Epi 1:1,000 per drug chart via nebulizer, MR x1</li> </ul> <p><b><u>Reassess following IM epinephrine. If no improvement in 2 minutes, consider:</u></b></p> <ul style="list-style-type: none"> <li>• Epi 1:1,000 per drug chart IM. MR x2 q5 minutes</li> </ul>

**Treatment Protocols**

**Date: 07/01/2023**

**Respiratory Distress or Failure - Pediatric**

**Policy #9170P**

<b>Pediatric ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• Monitor EKG</li> <li>• Establish IV/IO</li> <li>• Capnography</li> <li>• 12 Lead if cardiac source considered</li> </ul> <p><b><u>SUSPECTED BRONCHOSPASM</u></b></p> <ul style="list-style-type: none"> <li>• Albuterol weight based</li> <li>• Ipratropium weight based</li> <li>• Consider NIPPV PRN – See <b>NIPPV Procedure</b> (for adult sized pediatric patients only)</li> </ul> <p><b><u>CROUP / SUSPECTED CROUP</u></b></p> <ul style="list-style-type: none"> <li>• NS or Sterile Water 5 mL, via nebulizer mask, MR prn</li> </ul>	<ul style="list-style-type: none"> <li>• Monitor EKG</li> <li>• Establish IV/IO</li> <li>• Capnography</li> <li>• 12 Lead if cardiac source considered</li> </ul> <p><b><u>HYPOTENSION IF CARDIAC CAUSE NOT SUSPECTED</u></b></p> <ul style="list-style-type: none"> <li>• 10-20 mL/kg NS IV/IO bolus; titrated to age-appropriate systolic BP MR x1, if patient is without rales and there is no evidence of heart failure</li> </ul> <p><b><u>SUSPECTED BRONCHOSPASM</u></b></p> <ul style="list-style-type: none"> <li>• Albuterol weight based</li> <li>• Ipratropium weight based</li> <li>• Consider NIPPV PRN – See <b>NIPPV Procedure</b> (for adult sized pediatric patients only)</li> </ul> <p><b><u>CROUP / SUSPECTED CROUP</u></b></p> <ul style="list-style-type: none"> <li>• NS or Sterile Water 5 mL, via nebulizer, MR prn</li> </ul>
<b>Pediatric Base Hospital Orders</b>	
<p><b><u>EPIGLOTTITIS/ SUSPECTED EPIGLOTTITIS W/ STRIDOR</u></b></p> <ul style="list-style-type: none"> <li>• BHP – Epinephrine 1:1,000 weight based via nebulizer, monitor ECG during administration</li> </ul>	<p><b><u>EPIGLOTTITIS/ SUSPECTED EPIGLOTTITIS W/ STRIDOR</u></b></p> <ul style="list-style-type: none"> <li>• BHP – Epinephrine 1:1,000 weight based via nebulizer, monitor ECG during administration</li> </ul>
<b>Notes:</b>	
<ul style="list-style-type: none"> <li>• <b>Not all wheezing is from bronchospasm. A cardiac wheeze can occur from heart failure. If a pediatric patient has known cardiac history (congenital heart abnormality or Kawasaki’s disease for example) consider early Base Station contact and NIPPV.</b></li> <li>• <b>If a pediatric patient presents with stridor or significant upper airway noise, consider foreign body ingestion/aspiration as source of distress</b></li> <li>• <b>NIPPV can increase intrathoracic pressure and drop a patient’s blood pressure. Perform frequent BP rechecks, and do not use in profound or refractory hypotension</b></li> </ul>	

APPROVED:

Signature on File

Katherine Staats, M.D. FACEP

EMS Medical Director